

### **International Office**

#### **Everest Biotech Ltd**

Vector Laboratories, Inc. 6737 Mowry Ave Newark, CA 94560 United States

**Customer Service:** 

customerservice@vectorlabs.com

Technical Service:

technical@vectorlabs.com

Tel: +1 (800) 227-6666

www.everestbiotech.com

Research Use Only. Not for diagnostic or therapeutic use.

# EB09610 - Goat Anti-TPCN2 Antibody

Size: 100µg specific antibody in 200µl



# **Target Protein**

Principal Names: TPCN2, two pore segment channel 2, FLJ41094, SHEP10, TPC2,

two-pore calcium channel protein 2

Official Symbol: TPCN2

Accession Number(s): NP\_620714.2

Human GenelD(s): 219931

# **Immunogen**

Peptide with sequence C-EDELTERLSQHPH, from the C Terminus of the protein sequence according to NP\_620714.2.

Please note the peptide is available for sale.

# **Purification and Storage**

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin

Aliquot and store at -20°C. Minimize freezing and thawing.

## **Applications Tested**

Peptide ELISA: antibody detection limit dilution 1:16000.


**Western blot:** Preliminary experiments gave an approx 30kDa band in Human Brain (Cerebral Cortex), Kidney and Liver lysates after 0.5µg/ml antibody staining. Please note that currently we cannot find an explanation in the literature for the band we observe given the calculated size of 85.2kDa according to NP\_620714.2. The 30kDa band was successfully blocked by incubation with the immunizing peptide.

**IHC:** Paraffin embedded Human Pancreas and Colon. Recommended concentration: 3.75µg/ml.


### **Species Reactivity**

Tested: Human

Expected from sequence similarity: Human



EB09610 (3.75 $\mu$ g/ml) staining of paraffin embedded Human Pancreas. Steamed antigen retrieval with citrate buffer pH 6, AP-staining.



EB09610 (3.75 $\mu$ g/ml) staining of paraffin embedded Human Colon. Steamed antigen retrieval with citrate buffer pH 6, AP-staining.