

## UK Office

### Everest Biotech Ltd

Cherwell Innovation Centre  
77 Heyford Park  
Upper Heyford  
Oxfordshire  
OX25 5HD  
UK

Enquiries:

[info@everestbiotech.com](mailto:info@everestbiotech.com)

Sales:

[sales@everestbiotech.com](mailto:sales@everestbiotech.com)

Tech support:

[support@everestbiotech.com](mailto:support@everestbiotech.com)

Tel: +44 (0)1869 238326

[www.everestbiotech.com](http://www.everestbiotech.com)

**Research Use Only. Not for  
diagnostic or therapeutic use.**

## EB11012 - Goat Anti-Tph2 (mouse) Antibody

Size: 100µg specific antibody in 200µl



### Target Protein

**Principal Names:** Tph2, tryptophan hydroxylase 2, AU043594, MGC159133, Ntph, OTTMUSP00000022885, neuronal tryptophan hydroxylase, tryptophan 5-hydroxylase 2, tryptophan 5-monoxygenase 2, AU043594, MGC159133, neuronal tryptophan hydroxylase, Ntph, OTTMUSP00000022885, tryptophan 5-hydroxylase 2, tryptophan 5-monoxygenase 2, tryptophan hydroxylase 2, Tph2

**Official Symbol:** Tph2

**Accession Number(s):** NP\_775567.2

**Non-Human GeneID(s):** 216343 (mouse)

### Immunogen

Peptide with sequence C-SLTQNKAIKSEDK, from the internal region of the protein sequence according to NP\_775567.2.

Please note the [peptide](#) is available for sale.

### Purification and Storage

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

### Applications Tested

**Peptide ELISA:** antibody detection limit dilution 1:32000.

**Western blot:** Preliminary experiments gave an approx 85kDa band in Mouse fetal Brain lysates after 0.3µg/ml antibody staining. Please note that currently we cannot find an explanation in the literature for the band we observe given the calculated size of 55.9kDa according to NP\_775567.2. The 85kDa band was successfully blocked by incubation with the immunizing peptide.

**Immunofluorescence:** In free-floating samples of Mouse Brain stem staining of Raphe nuclei serotonergic neurons can be seen, that is consistent with observation by different antibodies. Recommended concentration, 0.5-2µg/ml.

### Species Reactivity

**Tested:** Mouse

**Expected from sequence similarity:** Mouse


### Specific Reference

**This antibody has been successfully used in IF on Mouse:**

Rahim RS, Meedeniya AC, Crane DI.

Central serotonergic neuron deficiency in a mouse model of Zellweger syndrome. Neuroscience. 2014 Aug 22;274:229-41.

PMID: 24881576



EB11012 (0.5µg/ml) staining of Mouse Brain stem in sagittal (left) and coronal (right) sections of Raphe nuclei.  
Detection Alexa 594. Data obtained from Prof. D Crane, Griffith Univeristy, Brisbane, Australia