



UK Office

Everest Biotech Ltd

Cherwell Innovation Centre
77 Heyford Park
Upper Heyford
Oxfordshire
OX25 5HD
UK

Enquiries:

info@everestbiotech.com

Sales:

sales@everestbiotech.com

Tech support:

support@everestbiotech.com

Tel: +44 (0)1869 238326

Fax: +44 (0)1869 238327

US Office

Everest Biotech c/o Abcore

405 Maple Street, Suite A106
Ramona,
CA 92065
USA

Inquiries:

info@everestbiotech.com

Sales:

usasales@everestbiotech.com

Tech support:

support@everestbiotech.com

Tel: 888-320-4628 (toll-free)

Fax: 888-841-9041

www.everestbiotech.com

**Research Use Only. Not for
diagnostic or therapeutic use.**

EB11012 - Goat Anti-Tph2 (mouse) Antibody

Size: 100µg specific antibody in 200µl



Target Protein

Principal Names: Tph2, tryptophan hydroxylase 2, AU043594, MGC159133, Ntph, OTTMUSP00000022885, neuronal tryptophan hydroxylase, tryptophan 5-hydroxylase 2, tryptophan 5-monooxygenase 2, AU043594, MGC159133, neuronal tryptophan hydroxylase, Ntph, OTTMUSP00000022885, tryptophan 5-hydroxylase 2, tryptophan 5-monooxygenase 2, tryptophan hydroxylase 2, Tph2

Official Symbol: Tph2

Accession Number(s): NP_775567.2

Non-Human GeneID(s): 216343 (mouse)

Immunogen

Peptide with sequence C-SLTQNKAIKSEDK, from the internal region of the protein sequence according to NP_775567.2.

Please note the [peptide](#) is available for sale.

Purification and Storage

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

Applications Tested

Peptide ELISA: antibody detection limit dilution 1:32000.

Western blot: Preliminary experiments gave an approx 85kDa band in Mouse fetal Brain lysates after 0.3µg/ml antibody staining. Please note that currently we cannot find an explanation in the literature for the band we observe given the calculated size of 55.9kDa according to NP_775567.2. The 85kDa band was successfully blocked by incubation with the immunizing peptide.

Immunofluorescence: In free-floating samples of Mouse Brain stem staining of Raphe nuclei serotonergic neurons can be seen, that is consistent with observation by different antibodies. Recommended concentration, 0.5-2µg/ml.

Species Reactivity

Tested: Mouse

Expected from sequence similarity: Mouse


Specific Reference

This antibody has been successfully used in IF on Mouse:

Rahim RS, Meedeniya AC, Crane DI.

Central serotonergic neuron deficiency in a mouse model of Zellweger syndrome. Neuroscience. 2014 Aug 22;274:229-41.

PMID: 24881576



EB11012 (0.5µg/ml) staining of Mouse Brain stem in sagittal (left) and coronal (right) sections of Raphe nuclei.
Detection Alexa 594. Data obtained from Prof. D Crane, Griffith Univeristy, Brisbane, Australia