



UK Office

Everest Biotech Ltd

Cherwell Innovation Centre
77 Heyford Park
Upper Heyford
Oxfordshire
OX25 5HD
UK

Enquiries:

info@everestbiotech.com

Sales:

sales@everestbiotech.com

Tech support:

support@everestbiotech.com

Tel: +44 (0)1869 238326

Fax: +44 (0)1869 238327

US Office

Everest Biotech c/o Abcore

405 Maple Street, Suite A106
Ramona,
CA 92065
USA

Inquiries:

info@everestbiotech.com

Sales:

usasales@everestbiotech.com

Tech support:

support@everestbiotech.com

Tel: 888-320-4628 (toll-free)

Fax: 888-841-9041

www.everestbiotech.com

**Research Use Only. Not for
diagnostic or therapeutic use.**

EB11081 - Goat Anti-Transglutaminase 2 (aa578-591) Antibody

Size: 100µg specific antibody in 200µl



Target Protein

Principal Names: C polypeptide, G-ALPHA-h, GNAH, protein-glutamine gamma-glutamyltransferase 2, protein-glutamine-gamma-glutamyltransferase, TG(C), TG2, TGase C, TGase H, TGase-2, TGase-H, TGC, tissue transglutaminase, transglutaminase 2 (C polypeptide, protein-glutamine-gamma-glutamyltransferase), transglutaminase C, transglutaminase H, transglutaminase-2, TGM2

Official Symbol: TGM2

Accession Number(s): NP_004604.2

Human GeneID(s): [7052](#)

Non-Human GeneID(s): 21817 (mouse), 56083 (rat)

Important Comments: This antibody is expected to recognize isoform a (NP_004604.2) only.

Immunogen

Peptide with sequence C-ERDLYLENPEIKIR, from the internal region of the protein sequence according to NP_004604.2.

Please note the [peptide](#) is available for sale.

Purification and Storage

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

Applications Tested

Peptide ELISA: antibody detection limit dilution 1:8000.

Western blot: Approx 80kDa band observed in Mouse and Rat Kidney lysates (calculated MW of 77.3kDa according to Human NP_004604.2 and 77.0kDa according to Mouse NP_033399.1). Recommended concentration: 1-3µg/ml.

Species Reactivity

Tested: Mouse, Rat

Expected from sequence similarity: Human, Mouse, Rat, Pig, Cow



EB11081 (1 μ g/ml) staining of Mouse Kidney lysate (35 μ g protein in RIPA buffer). Primary incubation was 1 hour.
Detected by chemiluminescence.