

International Office

Everest Biotech Ltd

Vector Laboratories, Inc.
6737 Mowry Ave
Newark, CA 94560
United States

Customer Service:

customerservice@vectorlabs.com

Technical Service:

technical@vectorlabs.com

Tel: +1 (800) 227-6666

www.everestbiotech.com

**Research Use Only. Not for
diagnostic or therapeutic use.**

EB05902 - Goat Anti-TRC8 Antibody

Size: 100µg specific antibody in 200µl



Target Protein

Principal Names: RNF139, ring finger protein 139, TRC8, HRCA1, MGC31961, RCA1, patched related protein translocated in renal cancer, multiple membrane spanning receptor TRC8

Official Symbol: RNF139

Accession Number(s): NP_009149.2

Human GeneID(s): [11236](#)

Immunogen

Peptide with sequence C-HTGAAAEFNDT, from the C Terminus of the protein sequence according to NP_009149.2.

Please note the [peptide](#) is available for sale.

Purification and Storage

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

Applications Tested

Peptide ELISA: antibody detection limit dilution 1:16000.


Western blot: Approx 75kDa band observed in Human Skeletal Muscle lysates (calculated MW of 76.0kDa according to NP_009149.2). Recommended concentration: 0.3-1µg/ml.

IHC: In paraffin embedded Human Placenta shows membranous staining in the cytotrophoblasts. Recommended concentration, 3-4µg/ml.


Species Reactivity

Tested: Human

Expected from sequence similarity: Human, Dog



EB05902 (1µg/ml) staining of Human Skeletal Muscle lysate (35µg protein in RIPA buffer). Primary incubation was 1 hour. Detected by chemiluminescence.



EB05902 (3.8µg/ml) staining of paraffin embedded Human Placenta. Steamed antigen retrieval with citrate buffer pH 6, AP-staining.