



UK Office

Everest Biotech Ltd

Cherwell Innovation Centre
77 Heyford Park
Upper Heyford
Oxfordshire
OX25 5HD
UK

Enquiries:

info@everestbiotech.com

Sales:

sales@everestbiotech.com

Tech support:

support@everestbiotech.com

Tel: +44 (0)1869 238326

Fax: +44 (0)1869 238327

US Office

Everest Biotech c/o Abcore

405 Maple Street, Suite A106
Ramona,
CA 92065
USA

Inquiries:

info@everestbiotech.com

Sales:

usasales@everestbiotech.com

Tech support:

support@everestbiotech.com

Tel: 888-320-4628 (toll-free)

Fax: 888-841-9041

www.everestbiotech.com

**Research Use Only. Not for
diagnostic or therapeutic use.**

EB05129 - Goat Anti-TRPM7 / LTRPC7 Antibody

Size: 100µg specific antibody in 200µl



Target Protein

Principal Names: TRPM7, transient receptor potential cation channel, subfamily M, member 7, CHAK, CHAK1, LTRPC7, FLJ20117, TRP-PLIK, LTRPC ion channel family member 7, homolog of mouse transient receptor potential-phospholipase C-interacting kinase CHaK hypothetical protein FLJ20117, FLJ25718

Official Symbol: TRPM7

Accession Number(s): NP_060142.3; NP_001288141.1

Human GeneID(s): [54822](#)

Immunogen

Peptide with sequence C-TKESESTNSVRLML, from the C Terminus of the protein sequence according to NP_060142.3; NP_001288141.1.

Please note the [peptide](#) is available for sale.

Purification and Storage

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

Applications Tested

Peptide ELISA: antibody detection limit dilution 1:8000.


IHC: In paraffin embedded Mouse Brain shows neuronal staining in the Hippocampus.
Recommended concentration: 4-6µg/ml.

Immunofluorescence: Strong expression of the protein seen in the cytoplasm and nuclei of U2OS cells. Recommended concentration: 10µg/ml.


Species Reactivity

Tested: Human, Mouse

Expected from sequence similarity: Human, Mouse, Rat, Dog, Pig



EB05129 (4 μ g/ml) staining of paraffin embedded Mouse Hippocampus. Steamed antigen retrieval with citrate buffer pH 6, HRP-staining. Similar results were obtained after antigen retrieval at pH9.



EB05129 Immunofluorescence analysis of paraformaldehyde fixed U2OS cells, permeabilized with 0.15% Triton. Primary incubation 1hr (10 μ g/ml) followed by Alexa Fluor 488 secondary antibody (2 μ g/ml), showing nuclear and cytoplasmic staining. Actin filaments were stained with phalloidin (red) and the nuclear stain is DAPI (blue). Negative control: Unimmunized goat IgG (10 μ g/ml) followed by Alexa Fluor 488 secondary antibody (2 μ g/ml).