

## **International Office**

#### Everest Biotech Ltd

Vector Laboratories, Inc. 6737 Mowry Ave Newark, CA 94560 United States

Customer Service: <u>customerservice@vectorlabs.com</u> Technical Service: <u>technical@vectorlabs.com</u>

Tel: +1 (800) 227-6666

www.everestbiotech.com

Research Use Only. Not for diagnostic or therapeutic use.

# EB08540 - Goat Anti-TXNDC4 Antibody

Size: 100µg specific antibody in 200µl



## **Target Protein**

**Principal Names:** TXNDC4, thioredoxin domain containing 4 (endoplasmic reticulum), ERP44, KIAA0573, OTTHUMP0000063799, endoplasmic reticulum resident protein 44

kDa Official Symbol: ERP44 Accession Number(s): NP\_055866.1 Human GeneID(s): 23071 Non-Human GeneID(s): 76299 (mouse), 298066 (rat)

#### Immunogen

Peptide with sequence C-EQKDSDNYRVFER, from the internal region of the protein sequence according to NP\_055866.1.

Please note the peptide is available for sale.

# **Purification and Storage**

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide. Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin. Aliquot and store at -20°C. Minimize freezing and thawing.

## Applications Tested

Peptide ELISA: antibody detection limit dilution 1:32000.

**Western blot:** Approx 47kDa band observed in lysates of cell line K562 (calculated MW of 47.0kDa according to NP\_055866.1). Recommended concentration: 0.2-0.6µg/ml. An additional band of unknown identity was also consistently observed at 22kDa. This band was successfully blocked by incubation with the immunising peptide. We would appreciate any feedback from people in the field - have any such results been reported with other antibodies/lysates? Have any further splice variants/modified forms been reported?

# **Species Reactivity**

Tested: Human Expected from sequence similarity: Human, Mouse, Rat, Cow

	250kDa 150kDa 100kDa 75kDa 50kDa
	37kDa
	25kDa
-	20kDa
	15kDa

EB08540 (0.2µg/ml) staining of K562 lysate (35µg protein in RIPA buffer). Primary incubation was 1 hour. Detected by chemiluminescence.