

### **International Office**

#### **Everest Biotech Ltd**

Vector Laboratories, Inc. 6737 Mowry Ave Newark, CA 94560 United States

**Customer Service:** 

customerservice@vectorlabs.com

Technical Service:

technical@vectorlabs.com

Tel: +1 (800) 227-6666

www.everestbiotech.com

Research Use Only. Not for diagnostic or therapeutic use.

# EB12639 - Goat Anti-U-PAR / CD87 Antibody

Size: 100µg specific antibody in 200µl



### **Target Protein**

**Principal Names:** PLAUR, plasminogen activator, urokinase receptor, CD87, U-PAR, UPAR, URKR, monocyte activation antigen Mo3, u-plasminogen activator receptor form 2, urokinase plasminogen activator surface receptor, urokinase-type plasminogen activator (uPA) receptor

Official Symbol: PLAUR

Accession Number(s): NP\_002650.1; NP\_001005376.1; NP\_001005377.1

Human GeneID(s): 5329

Important Comments: This antibody is expected to recognize all reported isoforms

(NP\_002650.1; NP\_001005376.1; NP\_001005377.1).

# **Immunogen**

Peptide with sequence C-RTGLKITSLTE , from the internal region of the protein sequence according to NP\_002650.1; NP\_001005376.1; NP\_001005377.1.

Please note the peptide is available for sale.

# **Purification and Storage**

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

## **Applications Tested**

Peptide ELISA: antibody detection limit dilution 1:128000.

**Western blot:** Approx 60kDa band observed in lysates of cell line U251 (calculated MW of 37.0kDa according to NP\_002650.1). The observed molecular weight corresponds to the glycosylated form. Recommended concentration: 1-3µg/ml.

### **Species Reactivity**

Tested: Human

Expected from sequence similarity: Human

250kDa 150kDa 100kDa 75kDa 50kDa 37kDa 25kDa 20kDa

EB12639 (1 $\mu$ g/ml) staining of U251 lysate (35 $\mu$ g protein in RIPA buffer). Primary incubation was 1 hour. Detected by chemiluminescence.