



UK Office

Everest Biotech Ltd

Cherwell Innovation Centre
77 Heyford Park
Upper Heyford
Oxfordshire
OX25 5HD
UK

Enquiries:

info@everestbiotech.com

Sales:

sales@everestbiotech.com

Tech support:

support@everestbiotech.com

Tel: +44 (0)1869 238326

Fax: +44 (0)1869 238327

US Office

Everest Biotech c/o Abcore

405 Maple Street, Suite A106
Ramona,
CA 92065
USA

Inquiries:

info@everestbiotech.com

Sales:

usasales@everestbiotech.com

Tech support:

support@everestbiotech.com

Tel: 888-320-4628 (toll-free)

Fax: 888-841-9041

www.everestbiotech.com

**Research Use Only. Not for
diagnostic or therapeutic use.**

EB05569 - Goat Anti-UBE2C / UBCH10 Antibody

Size: 100µg specific antibody in 200µl



Target Protein

Principal Names: UBE2C, UBCH10, dJ447F3.2, ubiquitin-conjugating enzyme E2C, ubiquitin-protein ligase C, ubiquitin carrier protein E2-C, cyclin-selective ubiquitin carrier protein, mitotic-specific ubiquitin-conjugating enzyme, RP3-447F3.1

Official Symbol: UBE2C

Accession Number(s): NP_008950.1; NP_861515.1; NP_861516.1; NP_861517.1; NP_861518.1; NP_001268670.1

Human GeneID(s): [11065](#)

Important Comments: This antibody is expected to recognise all reported isoforms. Variants NP_861517.1 and NP_861518.1 encode the same isoform.

Immunogen

Peptide with sequence C-QETYSKQVTSQEP, from the C Terminus of the protein sequence according to NP_008950.1; NP_861515.1; NP_861516.1; NP_861517.1; NP_861518.1; NP_001268670.1.

Please note the [peptide](#) is available for sale.

Purification and Storage

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

Applications Tested

Peptide ELISA: antibody detection limit dilution 1:32000.

Western blot: Approx.19KDa band observed in lysates of cell lines HeLa and HEK293 (calculated MW of 19.7kDa according to NP_008950.1). Recommended concentration: 1-3µg/ml. Primary incubation 1 hour at room temperature.

Immunofluorescence: Strong expression of the protein seen in the cytoplasm of U2OS and MCF7 cells. Recommended concentration: 10µg/ml.

Flow Cytometry: Flow cytometric analysis of HeLa cells. Recommended concentration: 10ug/ml.

Immunoprecipitation: 20KDa band precipitated from mitotic HeLa whole cell lysates using protein-G dynabeads.


Species Reactivity

Tested: Human


Expected from sequence similarity: Human




EB05569 (2 μ g/ml) staining of HEK293 (A) and HeLa (B) cell lysate (35 μ g protein in RIPA buffer). Detected by chemiluminescence.



EB05569 Immunofluorescence analysis of paraformaldehyde fixed U2OS cells, permeabilized with 0.15% Triton. Primary incubation 1hr (10ug/ml) followed by Alexa Fluor 488 secondary antibody (4ug/ml), showing cytoplasmic /Plasma Membrane staining. The nuclear stain is DAPI (blue). Negative control: Unimmunized goat IgG (10ug/ml) followed by Alexa Fluor 488 secondary antibody (4ug/ml).



EB05569 Immunofluorescence analysis of paraformaldehyde fixed MCF7 cells, permeabilized with 0.15% Triton. Primary incubation 1hr (10ug/ml) followed by Alexa Fluor 488 secondary antibody (4ug/ml), showing cytoplasmic staining. The nuclear stain is DAPI (blue). Negative control: Unimmunized goat IgG (10ug/ml) followed by Alexa Fluor 488 secondary antibody (4ug/ml).



EB05569 Flow cytometric analysis of paraformaldehyde fixed HeLa cells (blue line), permeabilized with 0.5% Triton. Primary incubation 1hr (10ug/ml) followed by Alexa Fluor 488 secondary antibody (2ug/ml). IgG control:

Unimmunized goat IgG (black line) followed by Alexa Fluor 488 secondary antibody.