

#### **International Office**

#### **Everest Biotech Ltd**

Vector Laboratories, Inc. 6737 Mowry Ave Newark, CA 94560 United States

**Customer Service:** 

customerservice@vectorlabs.com

Technical Service:

technical@vectorlabs.com

Tel: +1 (800) 227-6666

www.everestbiotech.com

Research Use Only. Not for diagnostic or therapeutic use.

# EB09611 - Goat Anti-UBR4 / ZUBR1 Antibody

Size: 100µg specific antibody in 200µl



## **Target Protein**

**Principal Names:** UBR4, ubiquitin protein ligase E3 component n-recognin 4, FLJ41863, KIAA0462, KIAA1307, RBAF600, ZUBR1, p600, retinoblastoma-associated factor 600, retinoblastoma-associated factor 600-like protein, zinc finger, UBR1 type 1

Official Symbol: UBR4

Accession Number(s): NP\_065816.2

Human GeneID(s): 23352

Non-Human GenelD(s): 69116 (mouse), 313658 (rat)

### **Immunogen**

Peptide with sequence C-NTTRATSREEKNLQ, from the C Terminus of the protein sequence according to NP\_065816.2.

Please note the peptide is available for sale.

# **Purification and Storage**

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

#### **Applications Tested**

Peptide ELISA: antibody detection limit dilution 1:128000.


**Western blot:** Not yet tested - our routinely used western blotting protocol does not allow detection of proteins as large as the calculated size of 574kDa according to NP\_065816.2. Therefore we cannot recommend an optimal concentration and the antibody is an aspiring product. We would appreciate any feedback from people in the field - have any results been reported with other antibodies/lysates? Have any further splice variants/modified forms been reported?

**IHC:** In paraffin embedded Spleen shows various types of subnuclear staining in select splenocytes and additional membrane plus cytoplasm staining elsewhere. Recommended concentration, 5-10µg/ml.

# **Species Reactivity**

Tested: Human

Expected from sequence similarity: Human, Mouse, Rat, Dog, Cow



EB09611 (5µg/ml) staining of paraffin embedded Human Spleen. Steamed antigen retrieval with citrate buffer pH 6, AP-staining.