

## UK Office

### Everest Biotech Ltd

Cherwell Innovation Centre  
77 Heyford Park  
Upper Heyford  
Oxfordshire  
OX25 5HD  
UK

Enquiries:

[info@everestbiotech.com](mailto:info@everestbiotech.com)

Sales:

[sales@everestbiotech.com](mailto:sales@everestbiotech.com)

Tech support:

[support@everestbiotech.com](mailto:support@everestbiotech.com)

Tel: +44 (0)1869 238326

[www.everestbiotech.com](http://www.everestbiotech.com)

**Research Use Only. Not for  
diagnostic or therapeutic use.**

## EB05416 - Goat Anti-UEV2V2 / MMS2 Antibody

Size: 100µg specific antibody in 200µl



### Target Protein

**Principal Names:** UBE2V2, MMS2, ubiquitin-conjugating enzyme E2 variant 2, UEV2, EDPF1, UEV-2, DDVIT1, EDAF-1, EDPF-1, DDVit-1, 1 alpha,25-dihydroxyvitamin D3-inducible, enterocyte differentiation promoting factor, methyl methanesulfonate sensitive 2, *S. cerevisiae*, homolog of, ubiquitin-conjugating enzyme E2v2

**Official Symbol:** UBE2V2

**Accession Number(s):** NP\_003341.1; NP\_954673.1

**Human GeneID(s):** [7336](#)

**Non-Human GeneID(s):** 70620 (mouse)

**Important Comments:** This antibody is also expected to recognize the readthrough transcript product KUA-UEV (NP\_954673.1, GeneID 387522). Also cross-reactivity is expected with UEV2V1 (GeneID 7335).

### Immunogen

Peptide with sequence AVSTGVKVP RNF-C, from the N Terminus of the protein sequence according to NP\_003341.1; NP\_954673.1.

Please note the [peptide](#) is available for sale.

### Purification and Storage

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

### Applications Tested

**Peptide ELISA:** antibody detection limit dilution 1:32000.

**Western blot:** Approx 20kDa band seen in human brain lysate. Recommended for use at 0.2-1µg/ml.

### Species Reactivity

**Tested:** Human

**Expected from sequence similarity:** Human, Mouse, Rat, Dog, Cow



EB05416 staining (0.5 $\mu$ g/ml) of human brain lysate (RIPA buffer, 30 $\mu$ g total protein per lane). Primary incubated for 12 hour. Detected by western blot using chemiluminescence.