



## International Office

### Everest Biotech Ltd

Vector Laboratories, Inc.  
6737 Mowry Ave  
Newark, CA 94560  
United States

Customer Service:

[customerservice@vectorlabs.com](mailto:customerservice@vectorlabs.com)

Technical Service:

[technical@vectorlabs.com](mailto:technical@vectorlabs.com)

Tel: +1 (800) 227-6666

[www.everestbiotech.com](http://www.everestbiotech.com)

**Research Use Only. Not for  
diagnostic or therapeutic use.**

## EB11647 - Goat Anti-Uromodulin (aa525-536) Antibody

Size: 100µg specific antibody in 200µl



### Target Protein

**Principal Names:** ADMCKD2, FJHN, HNFJ, HNFJ1, MCKD2, OTTHUMP00000162212, OTTHUMP00000162213, Tamm-Horsfall glycoprotein, tamm-Horsfall urinary glycoprotein, THGP, THP, UMOD, uromodulin, uromodulin (uromucoid, Tamm-Horsfall glycoprotein), uromucoid

**Official Symbol:** UMOD

**Accession Number(s):** NP\_003352.2

**Human GeneID(s):** [7369](#)

**Important Comments:** Reported variants represent identical protein: NP\_001008390.1, NP\_003352.2

### Immunogen

Peptide with sequence DRCPHTRDSTIQ, from the internal region of the protein sequence according to NP\_003352.2.

Please note the [peptide](#) is available for sale.

### Purification and Storage

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

### Applications Tested


**Peptide ELISA:** antibody detection limit dilution 1:64000.

**Immunofluorescence:** Strong expression of the protein seen in the Plasma membrane and Golgi of HeLa cells. Recommended concentration: 10µg/ml.

### Species Reactivity

**Tested:** Human

**Expected from sequence similarity:** Human



EB11647 Immunofluorescence analysis of paraformaldehyde fixed HeLa cells, permeabilized with 0.15% Triton. Primary incubation 1hr (10ug/ml) followed by Alexa Fluor 488 secondary antibody (2ug/ml), showing membrane and Golgi staining. The nuclear stain is DAPI (blue). Negative control: Unimmunized goat IgG (10ug/ml) followed by Alexa Fluor 488 secondary antibody (2ug/ml).