

#### **International Office**

#### **Everest Biotech Ltd**

Vector Laboratories, Inc. 6737 Mowry Ave Newark, CA 94560 United States

**Customer Service:** 

customerservice@vectorlabs.com

Technical Service:

technical@vectorlabs.com

Tel: +1 (800) 227-6666

www.everestbiotech.com

Research Use Only. Not for diagnostic or therapeutic use.

# EB11496 - Goat Anti-USP6 (aa142-155) Antibody

Size: 100µg specific antibody in 200µl



### **Target Protein**

**Principal Names:** deubiquitinating enzyme 6, HRP1, hyperpolymorphic gene 1, OTTHUMP00000125301, proto-oncogene TRE-2, TRE17, TRE2, Tre-2, tre-2 oncogene, ubiquitin carboxyl-terminal hydrolase 6, ubiquitin specific peptidase 6-, ubiquitin specific protease 6 (Tre-2 oncogene), ubiquitin thioesterase 6, ubiquitin thiolesterase 6, ubiquitin thiolesterase 0, ubiquitin-specific protease USP6,

ubiquitin-specific-processing protease 6, USP6, USP6-short

Official Symbol: USP6

Accession Number(s): NP\_004496.2

Human GenelD(s): 9098

Non-Human GenelD(s): 237898 (mouse)

Important Comments: This antibody is expected NOT to cross react to USP32 or to the

TBC1D3 proteins.

#### **Immunogen**

Peptide with sequence C-HHIDLDVRTTLRNH, from the internal region (near N Terminus) of the protein sequence according to NP\_004496.2.

Please note the peptide is available for sale.

### **Purification and Storage**

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

### **Applications Tested**

Peptide ELISA: antibody detection limit dilution 1:4000.

**Western blot:** Preliminary experiments in Human Testis, Prostate and Placenta lysates gave no specific signal but low background at antibody concentration up to 1µg/ml.


**Immunofluorescence:** Strong expression of the protein seen in the cytoplasm of A431 and U2OS cells. Recommended concentration: 10µg/ml.

**Flow Cytometry:** Flow cytometric analysis of A431 cells. Recommended concentration: 10ug/ml.


## **Species Reactivity**

Tested: Human


Expected from sequence similarity: Human



EB11496 Immunofluorescence analysis of paraformaldehyde fixed A431 cells, permeabilized with 0.15% Triton. Primary incubation 1hr (10ug/ml) followed by Alexa Fluor 488 secondary antibody (2ug/ml), showing cytoplasmic/vesicle staining. The nuclear stain is DAPI (blue). Negative control: Unimmunized goat IgG (10ug/ml) followed by Alexa Fluor 488 secondary antibody (2ug/ml).



EB11496 Immunofluorescence analysis of paraformaldehyde fixed U2OS cells, permeabilized with 0.15% Triton. Primary incubation 1hr (10ug/ml) followed by Alexa Fluor 488 secondary antibody (2ug/ml), showing cytoplasmic staining. The nuclear stain is DAPI (blue). Negative control: Unimmunized goat IgG (10ug/ml) followed by Alexa Fluor 488 secondary antibody (2ug/ml).



EB11496 Flow cytometric analysis of paraformaldehyde fixed A431 cells (blue line), permeabilized with 0.5% Triton. Primary incubation overnight (10ug/ml) followed by Alexa Fluor 488 secondary antibody (1ug/ml). IgG control: Unimmunized goat IgG (black line) followed by Alexa Fluor 488 secondary antibody.