

#### **International Office**

#### **Everest Biotech Ltd**

Vector Laboratories, Inc. 6737 Mowry Ave Newark, CA 94560 United States

**Customer Service:** 

customerservice@vectorlabs.com

Technical Service:

technical@vectorlabs.com

Tel: +1 (800) 227-6666

www.everestbiotech.com

Research Use Only. Not for diagnostic or therapeutic use.

# EB05974 - Goat Anti-USP6 / TRE2 / TRE17 Antibody

Size: 100µg specific antibody in 200µl



## **Target Protein**

**Principal Names:** USP6, ubiquitin specific protease 6 (Tre-2 oncogene), HRP1, TRE2, Tre2, TRE17, Tre-2, tre-2 oncogene, hyperpolymorphic gene 1, USP6-short, ubiquitin specific peptidase 6-, ubiquitin specific protease 6, ubiquitin specific peptidase 6 (Tre-2 oncogene), ubiquitin-specific protease USP6

Official Symbol: USP6

Accession Number(s): NP\_004496.1

Human GeneID(s): 9098

#### **Immunogen**

Peptide with sequence KISPLHHLQMECSP, from the C Terminus of the protein sequence according to NP\_004496.1.

Please note the peptide is available for sale.

### **Purification and Storage**

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

#### **Applications Tested**


Peptide ELISA: antibody detection limit dilution 1:16000.

Western blot: Approx 90kDa band observed in Human Placenta lysates (calculated MW of 89.5kDa according to NP\_004496.1). Recommended concentration: 1-3μg/ml. An additional band of 50kDa was consistently observed, however this band was not blocked by the immunizing peptide and it is therefore a non-specific signal. We call for caution when used for other assays than Western blot.

#### **Species Reactivity**

Tested: Human

Expected from sequence similarity: Human



EB05974 ( $2\mu g/ml$ ) staining of Human Placenta lysate ( $35\mu g$  protein in RIPA buffer) with (B) and without (A) blocking with the immunizing peptide. Primary incubation was 1 hour. Detected by chemiluminescence.