



## International Office

### Everest Biotech Ltd

Vector Laboratories, Inc.  
6737 Mowry Ave  
Newark, CA 94560  
United States

Customer Service:

[customerservice@vectorlabs.com](mailto:customerservice@vectorlabs.com)

Technical Service:

[technical@vectorlabs.com](mailto:technical@vectorlabs.com)

Tel: +1 (800) 227-6666

[www.everestbiotech.com](http://www.everestbiotech.com)

**Research Use Only. Not for  
diagnostic or therapeutic use.**

## EB06803 - Goat Anti-VDR Antibody

Size: 100µg specific antibody in 200µl



### Target Protein

**Principal Names:** VDR, vitamin D (1,25- dihydroxyvitamin D3) receptor, HGNC:12679, NR1H1, vitamin D (1,25-dihydroxyvitamin D3) receptor

**Official Symbol:** VDR

**Accession Number(s):** NP\_000367.1; NP\_001017535.1;  
NP\_001351014.1;NP\_001017536.1

**Human GeneID(s):** [7421](#)

**Non-Human GeneID(s):** 22337 (mouse), 24873 (rat)

### Immunogen

Peptide with sequence CGNQDYKYRVSD, from the internal region of the protein sequence according to NP\_000367.1; NP\_001017535.1;  
NP\_001351014.1;NP\_001017536.1.

Please note the [peptide](#) is available for sale.

### Purification and Storage

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

### Applications Tested

**Peptide ELISA:** antibody detection limit dilution 1:16000.


**Western blot:** Approx 55kDa band observed in Human Placenta and Duodenum lysates and approx. 55-60kDa in lysates of cell line U937 (calculated MW of 55.2kDa according to NP\_001351014.1). Recommended concentration: 2-3µg/ml. Primary incubation 1 hour at room temperature.

**Flow Cytometry:** Flow cytometric analysis of U937 cells. Recommended concentration: 10ug/ml.


### Species Reactivity

**Tested:** Human


**Expected from sequence similarity:** Human, Mouse, Rat, Dog



EB06803 (3 $\mu$ g/ml) staining of U937 cell lysate (35 $\mu$ g protein in RIPA buffer. Detected by chemiluminescence.



EB06803 (3 $\mu$ g/ml) staining of Human Placenta (A) and (2 $\mu$ g/ml) Duodenum (B) lysate (35 $\mu$ g protein in RIPA buffer. Detected by chemiluminescence.



EB06803 Flow cytometric analysis of paraformaldehyde fixed U937 cells (blue line), permeabilized with 0.5% Triton. Primary incubation 1hr (10ug/ml) followed by Alexa Fluor 488 secondary antibody (1ug/ml). IgG control: Unimmunized goat IgG (black line) followed by Alexa Fluor 488 secondary antibody.