

#### **International Office**

#### **Everest Biotech Ltd**

Vector Laboratories, Inc. 6737 Mowry Ave Newark, CA 94560 United States

**Customer Service:** 

customerservice@vectorlabs.com

Technical Service:

technical@vectorlabs.com

Tel: +1 (800) 227-6666

www.everestbiotech.com

Research Use Only. Not for diagnostic or therapeutic use.

# EB07333-T - Goat Anti-VHL Antibody - Trial

Size: 20µg specific antibody in 40µl



## **Target Protein**

**Principal Names:** VHL, von Hippel-Lindau tumor suppressor, E3 ubiquitin protein ligase, HRCA1, RCA1, VHL1, elongin binding protein, protein G7, pVHL

Official Symbol: VHL

Accession Number(s): NP\_000542.1; NP\_937799.1

Human GeneID(s): 7428

Non-Human GenelD(s): 22346 (mouse), 24874 (rat)

Important Comments: This antibody is expected to recognise both reported isoforms

(NP\_000542.1 and NP\_937799.1).

# Immunogen

Peptide with sequence C-RSLVKPENYRRLD, from the internal region of the protein sequence according to NP\_000542.1; NP\_937799.1.

Please note the peptide is available for sale.

## **Purification and Storage**

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

# **Applications Tested**

Peptide ELISA: antibody detection limit dilution 1:128000.

**Western blot:** Approx 17kDa band observed in Human Ovary and Testis lysates and also in lysates of Mouse and Rat Testis and Brain and of all human and mouse cell lines tested (calculated MW of 19.7kDa according to NP\_937799.1). Recommended concentration:

0.01-0.03µg/ml. Primary incubation was 1 hour.

IHC: Paraffin embedded Human Kidney. Recommended concentration: 5µg/ml.


## **Species Reactivity**

Tested: Human, Mouse, Rat

Expected from sequence similarity: Human, Mouse, Rat, Dog



EB07333 ( $0.01\mu g/ml$ ) staining of Human Ovary lysate ( $35\mu g$  protein in RIPA buffer). Detected by chemiluminescence.



EB07333 (5μg/ml) staining of paraffin embedded Human Kidney. Steamed antigen retrieval with citrate buffer pH 6, AP-staining.