

UK Office

Everest Biotech Ltd

Cherwell Innovation Centre 77 Heyford Park Upper Heyford Oxfordshire OX25 5HD

UK

Enquiries:

info@everestbiotech.com

Sales:

sales@everestbiotech.com

Tech support:

support@everestbiotech.com

Tel: +44 (0)1869 238326 Fax: +44 (0)1869 238327

US Office

Everest Biotech c/o Abcore

405 Maple Street, Suite A106 Ramona, CA 92065

USA

Inquiries:

info@everestbiotech.com

Sales:

usasales@everestbiotech.com

Tech support:

support@everestbiotech.com

Tel: 888-320-4628 (toll-free)

Fax: 888-841-9041

www.everestbiotech.com

Research Use Only. Not for diagnostic or therapeutic use.

EB07333-B - Goat Anti-VHL, Biotinylated Antibody

Size: 100µg specific antibody in 200µl

libody

Target Protein

Principal Names: VHL, von Hippel-Lindau tumor suppressor, E3 ubiquitin protein ligase, HRCA1, RCA1, VHL1, pVHL, elongin binding protein, protein G7

Official Symbol: VHL

Accession Number(s): NP_000542.1; NP_937799.1; XP_011532380.1

Human GeneID(s): 7428

Non-Human GenelD(s): 22346 (mouse), 24874 (rat)

Important Comments: This antibody is expected to recognise both reported isoforms

(NP_000542.1 and NP_937799.1).

Immunogen

Peptide with sequence C-RSLVKPENYRRLD., from the internal region of the protein sequence according to NP_000542.1; NP_937799.1; XP_011532380.1.

Please note the peptide is available for sale.

Purification and Storage

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin

Aliquot and store at -20°C. Minimize freezing and thawing.

Applications Tested

Peptide ELISA: antibody detection limit dilution 1:128000.

Western blot: Approx 17kDa band observed in Human Ovary lysates (calculated MW of 19.7kDa according to NP_937799.1). See non-biotinylated parental product's datasheet for further QC data. Recommended concentration: 0.1-0.3μg/ml.

Species Reactivity

Tested: Human, Mouse, Rat

Expected from sequence similarity: Human



Biotinylated EB07333 (0.3µg/ml) staining of Human Ovary lysate (35µg protein in RIPA buffer), exactly mirroring its parental non-biotinylated product. Primary incubation was 1 hour. Detected by chemiluminescence, using streptavidin-HRP and using NAP blocker as a substitute for skimmed milk.