

### **UK Office**

#### **Everest Biotech Ltd**

Cherwell Innovation Centre

77 Heyford Park Upper Heyford Oxfordshire OX25 5HD

UK

**Enquiries:** 

info@everestbiotech.com

Sales:

sales@everestbiotech.com

Tech support:

support@everestbiotech.com

Tel: +44 (0)1869 238326 Fax: +44 (0)1869 238327

#### **US Office**

### **Everest Biotech c/o Abcore**

405 Maple Street, Suite A106

Ramona, CA 92065 USA

Inquiries:

info@everestbiotech.com

Sales:

 $\underline{usasales@everest biotech.com}$ 

Tech support:

support@everestbiotech.com

Tel: 888-320-4628 (toll-free)

Fax: 888-841-9041

www.everestbiotech.com

Research Use Only. Not for diagnostic or therapeutic use.

# EB09860 - Goat Anti-VLCAD Antibody

Size: 100µg specific antibody in 200µl



## **Target Protein**

Principal Names: ACAD6, acyl-Coenzyme A dehydrogenase, very long chain, LCACD,

VLCAD, ACADVL

Official Symbol: ACADVL

Accession Number(s): NP\_000009.1; NP\_001029031.1

Human GeneID(s): 37

**Important Comments:** This antibody is expected to recognize both reported isoforms

(NP\_000009.1; NP\_001029031.1).

## **Immunogen**

Peptide with sequence DKSDSHPSDALTRK-C, from the N-Terminus (near) of the protein sequence according to NP\_000009.1; NP\_001029031.1.

Please note the peptide is available for sale.

## **Purification and Storage**

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

## **Applications Tested**

Peptide ELISA: antibody detection limit dilution 1:2000.

Western blot: Approx 68kDa band observed in Human Heart lysates (calculated MW of 68.1kDa according to NP\_001029031.1). Recommended concentration: 1-3μg/ml.

## **Species Reactivity**

Tested: Human

Expected from sequence similarity: Human

250kDa 150kDa 100kDa 75kDa 50kDa 37kDa 25kDa 20kDa

EB09860 (2μg/ml) staining of Human Heart lysate (35μg protein in RIPA buffer). Primary incubation was 1 hour.

Detected by chemiluminescence.