

## UK Office

### Everest Biotech Ltd

Cherwell Innovation Centre  
77 Heyford Park  
Upper Heyford  
Oxfordshire  
OX25 5HD  
UK

Enquiries:

[info@everestbiotech.com](mailto:info@everestbiotech.com)

Sales:

[sales@everestbiotech.com](mailto:sales@everestbiotech.com)

Tech support:

[support@everestbiotech.com](mailto:support@everestbiotech.com)

Tel: +44 (0)1869 238326

[www.everestbiotech.com](http://www.everestbiotech.com)

**Research Use Only. Not for  
diagnostic or therapeutic use.**

## EB06828 - Goat Anti-VPS11 Antibody

Size: 100µg specific antibody in 200µl



### Target Protein

**Principal Names:** VPS11, vacuolar protein sorting 11 (yeast), HGNC:14583, END1, PEP5, RNF108, hVPS11, vacuolar protein sorting 11 (yeast homolog), vacuolar protein sorting protein 11, vacuolar protein sorting 11 homolog (S. cerevisiae)

**Official Symbol:** VPS11

**Accession Number(s):** NP\_068375.3

**Human GeneID(s):** [55823](#)

**Non-Human GeneID(s):** 71732 (mouse)

### Immunogen

Peptide with sequence C-HALQEKDTQTKLE, from the internal region of the protein sequence according to NP\_068375.3.

Please note the [peptide](#) is available for sale.

### Purification and Storage

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

### Applications Tested

**Peptide ELISA:** antibody detection limit dilution 1:32000.

**Western blot:** Preliminary experiments gave an approx 75kDa band in human duodenum, colon and placenta lysates after 0.3µg/ml antibody staining. Please note that currently we cannot find an explanation in the literature for the band we observe given the calculated size of 108kDa according to NP\_068375.3. The 75kDa band was successfully blocked by incubation with the immunizing peptide.


**IHC:** In paraffin embedded Human Kidney shows brushborder staining in PCT cells and Human Pancreas shows cytoplasm staining of beta cells in islets of Langerhans.

Recommended concentration, 3-6µg/ml.


### Species Reactivity

**Tested:** Human

**Expected from sequence similarity:** Human, Mouse, Rat, Dog, Cow



EB06828 (3.8µg/ml) staining of paraffin embedded Human Kidney-. Steamed antigen retrieval with citrate buffer pH 6, AP-staining.



EB06828 (4µg/ml) staining of paraffin embedded Human Pancreas. Steamed antigen retrieval with Tris/EDTA buffer pH 9, HRP-staining.