

#### **International Office**

#### **Everest Biotech Ltd**

Vector Laboratories, Inc. 6737 Mowry Ave Newark, CA 94560 United States

**Customer Service:** 

customerservice@vectorlabs.com

Technical Service:

technical@vectorlabs.com

Tel: +1 (800) 227-6666

www.everestbiotech.com

Research Use Only. Not for diagnostic or therapeutic use.

# EB06257 - Goat Anti-VPS29 (C Terminus) Antibody

Size: 100µg specific antibody in 200µl

#### **Target Protein**

**Principal Names:** VPS29, DC7, DC15, PEP11, FLJ20492, DKFZp564F0223, vacuolar protein sorting 29 (yeast), x 007 protein, retromer protein, vacuolar sorting protein VPS29/PEP11, vacuolar protein sorting 29 (yeast homolog), vacuolar protein sorting 29 homolog (S. cerevisiae), vacuolar protein sorting 29

Official Symbol: VPS29

Accession Number(s): NP\_057310.1; NP\_476528.1

Human GeneID(s): 51699

Non-Human GenelD(s): 56433 (mouse)

Important Comments: This antibody is expected to recognise both human isoforms

(represented by NP\_057310.1; NP\_476528.1).

### **Immunogen**

Peptide with sequence C-DDVKVERIEYKKP, from the C Terminus of the protein sequence according to NP\_057310.1; NP\_476528.1.

Please note the peptide is available for sale.

# **Purification and Storage**

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

#### **Applications Tested**

Peptide ELISA: antibody detection limit dilution 1:16000.

**Western blot:** Approx 20+21kDa bands observed in Human Spleen lysates (calculated MW of 20.5kDa according to NP\_057310.1 and 20.9kDa according to NP\_476528.1). Recommended concentration: 0.1-0.3µg/ml. Primary incubated for 1 hour.

# **Species Reactivity**

Tested: Human

Expected from sequence similarity: Human, Mouse, Rat, Dog, Cow

250kDa 150kDa 100kDa 75kDa 50kDa 37kDa 25kDa 20kDa 15kDa

EB06257 staining (0.1µg/ml) of Human Spleen lysate (RIPA buffer, 35µg total protein per lane). Detected by chemiluminescence.