



## UK Office

### Everest Biotech Ltd

Cherwell Innovation Centre  
77 Heyford Park  
Upper Heyford  
Oxfordshire  
OX25 5HD  
UK

Enquiries:

[info@everestbiotech.com](mailto:info@everestbiotech.com)

Sales:

[sales@everestbiotech.com](mailto:sales@everestbiotech.com)

Tech support:

[support@everestbiotech.com](mailto:support@everestbiotech.com)

Tel: +44 (0)1869 238326

[www.everestbiotech.com](http://www.everestbiotech.com)

**Research Use Only. Not for  
diagnostic or therapeutic use.**

## EB09545 - Goat Anti-VRK2 Antibody

Size: 100µg specific antibody in 200µl



### Target Protein

**Principal Names:** VRK2, vaccinia related kinase 2, vaccinia virus B1R-related kinase 2, vaccinia-related kinase-2

**Official Symbol:** VRK2

**Accession Number(s):** NP\_001129499.1; NP\_006287.2; NP\_001123954.1; NP\_001123955.1

**Human GeneID(s):** [7444](#)

**Important Comments:** This antibody is expected to recognize all isoforms (NP\_001123952.1, NP\_001123954.1, NP\_001123955.1, NP\_001129499.1). Reported variants represent identical protein: NP\_001123953.1, NP\_006287.2.

### Immunogen

Peptide with sequence C-PNSQKVDSQKAATK, from the internal region of the protein sequence according to NP\_001129499.1; NP\_006287.2; NP\_001123954.1; NP\_001123955.1.

Please note the [peptide](#) is available for sale.

### Purification and Storage

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

### Applications Tested

**Peptide ELISA:** antibody detection limit dilution 1:4000.

**Western blot:** Approx 60kDa band observed in nuclear lysates of cell line K562 (calculated MW of 58.1kDa according to NP\_001123954.1). Recommended concentration: 1-3µg/ml.

### Species Reactivity

**Tested:** Human

**Expected from sequence similarity:** Human, Dog



EB09545 (2 $\mu$ g/ml) staining of K562 nuclear lysate (35 $\mu$ g protein in RIPA buffer). Primary incubation was 1 hour.  
Detected by chemiluminescence.