

UK Office

Everest Biotech Ltd

Cherwell Innovation Centre 77 Heyford Park Upper Heyford

Oxfordshire OX25 5HD

UK

Enquiries:

info@everestbiotech.com

Sales:

sales@everestbiotech.com

Tech support:

support@everestbiotech.com

Tel: +44 (0)1869 238326 Fax: +44 (0)1869 238327

US Office

Everest Biotech c/o Abcore

405 Maple Street, Suite A106

Ramona, CA 92065 USA

Inquiries:

info@everestbiotech.com

Sales:

usasales@everestbiotech.com

Tech support:

support@everestbiotech.com

Tel: 888-320-4628 (toll-free)

Fax: 888-841-9041

www.everestbiotech.com

Research Use Only. Not for diagnostic or therapeutic use.

EB07215 - Goat Anti-VTI1B Antibody

Size: 100µg specific antibody in 200µl



Target Protein

Principal Names: VTI1B, vesicle transport through interaction with t-SNAREs homolog 1B (yeast), HGNC:17793, VTI1, VTI1L, VTI2, v-SNARE, vesicle transport through interaction with t-SNAREs 1B, vesicle-associated soluble NSF attachment protein

receptor
Official Symbol: VTI1B

Accession Number(s): NP_006361.1

Human GeneID(s): 10490

Non-Human GeneID(s): 53612 (mouse)

Immunogen

Peptide with sequence C-RDFDEKQQEANET, from the internal region of the protein sequence according to NP_006361.1.

Please note the peptide is available for sale.

Purification and Storage

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin

Aliquot and store at -20°C. Minimize freezing and thawing.

Applications Tested

Peptide ELISA: antibody detection limit dilution 1:32000.

Western blot: Approx 26-28kDa band observed in Human Duodenum, Kidney and Testes

lysates (calculated MW of 26.7kDa according to NP_006361.1). Recommended

concentration: 1-3µg/ml.

Species Reactivity

Tested: Human

Expected from sequence similarity: Human, Mouse, Rat, Dog

250kDa 150kDa 100kDa 75kDa 50kDa 37kDa 25kDa 20kDa 15kDa

EB07215 (1μg/ml) staining of Human Duodenum lysate (35μg protein in RIPA buffer). Primary incubation was 1 hour. Detected by chemiluminescence.