



International Office

Everest Biotech Ltd

Vector Laboratories, Inc.
6737 Mowry Ave
Newark, CA 94560
United States

Customer Service:

customerservice@vectorlabs.com

Technical Service:

technical@vectorlabs.com

Tel: +1 (800) 227-6666

www.everestbiotech.com

**Research Use Only. Not for
diagnostic or therapeutic use.**

EB05374 - Goat Anti-WNT4 Antibody

Size: 100µg specific antibody in 200µl



Target Protein

Principal Names: WNT4, wingless-type MMTV integration site family, member 4, WNT-4, WNT-4 protein precursor, RP1-224A6.7, SERKAL, WNT-4 protein

Official Symbol: WNT4

Accession Number(s): NP_110388.2

Human GeneID(s): [54361](#)

Non-Human GeneID(s): 22417 (mouse)

Immunogen

Peptide with sequence C-SNWLYLAKLSSVGS, from the internal region of the protein sequence according to NP_110388.2.

Please note the [peptide](#) is available for sale.

Purification and Storage

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

Applications Tested

Peptide ELISA: antibody detection limit dilution 1:4000.


Western blot: Approx 40+35kDa band observed in lysates of cell line Jurkat (calculated MW of 39.1kDa according to NP_110388.2). Recommended concentration: 1-3µg/ml. Primary incubation 1 hour at room temperature. **Negative Control:** U937 cell lysate.

Immunofluorescence: Strong expression of the protein seen in the Vesicle of HeLa and MCF7 cells. Recommended concentration: 10µg/ml.


Species Reactivity

Tested: Human


Expected from sequence similarity: Human, Mouse, Rat



EB05374 (1 μ g/ml) staining of Jurkat (A) and negative control U937 (B) cell lysate (35 μ g protein in RIPA buffer).
Detected by chemiluminescence.



EB05374 Immunofluorescence analysis of paraformaldehyde fixed HeLa cells, permeabilized with 0.15% Triton.
Primary incubation 1hr (10 μ g/ml) followed by Alexa Fluor 488 secondary antibody (4 μ g/ml), showing vesicle staining. The nuclear stain is DAPI (blue). Negative control: Unimmunized goat IgG (10 μ g/ml) followed by Alexa Fluor 488 secondary antibody (4 μ g/ml).



EB05374 Immunofluorescence analysis of paraformaldehyde fixed MCF7 cells, permeabilized with 0.15% Triton. Primary incubation 1hr (10ug/ml) followed by Alexa Fluor 488 secondary antibody (4ug/ml), showing vesicle staining. The nuclear stain is DAPI (blue). Negative control: Unimmunized goat IgG (10ug/ml) followed by Alexa Fluor 488 secondary antibody (4ug/ml).