

## **UK Office**

#### **Everest Biotech Ltd**

Cherwell Innovation Centre 77 Heyford Park Upper Heyford Oxfordshire OX25 5HD

UK

**Enquiries:** 

info@everestbiotech.com

Sales:

sales@everestbiotech.com

Tech support:

support@everestbiotech.com

Tel: +44 (0)1869 238326 Fax: +44 (0)1869 238327

### **US Office**

#### **Everest Biotech c/o Abcore**

405 Maple Street, Suite A106

Ramona, CA 92065 USA

Inquiries:

info@everestbiotech.com

Sales:

 $\underline{usasales@everest biotech.com}$ 

Tech support:

 $\underline{support@everest biotech.com}$ 

Tel: 888-320-4628 (toll-free)

Fax: 888-841-9041

www.everestbiotech.com

Research Use Only. Not for diagnostic or therapeutic use.

# EB05062 - Goat Anti-XAB2 / HCNP Antibody

Size: 100µg specific antibody in 200µl



## **Target Protein**

Principal Names: XAB2, HCNP, XPA binding protein 2, HCRN, crn-related protein kim1,

DKFZp762C1015, NTC90, SYF1, SYF1 homolog, RNA splicing factor

Official Symbol: XAB2

Accession Number(s): NP\_064581.2

Human GenelD(s): 56949

### **Immunogen**

Peptide with sequence C-SVPAAVFGSLKED, from the C Terminus of the protein sequence according to NP\_064581.2.

Please note the peptide is available for sale.

## **Purification and Storage**

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

## **Applications Tested**

Peptide ELISA: antibody detection limit dilution 1:8000.

**Western blot:** Approx 90kDa band observed in human fibroblast A431 lysates (calculated MW of 100kDa according to NP\_064581.2). Recommended concentration: 0.3-1µg/ml. **IHC:** In paraffin embedded Human Brain (Cerebral Cortex) shows nuclear staining.

Recommended concentration, 3-4µg/ml.


## **Species Reactivity**

Tested: Human

Expected from sequence similarity: Human, Mouse, Rat, Dog, Cow



EB05062 staining (0.3μg/ml) of A431 lysate (RIPA buffer, 35μg total protein per lane). Primary incubated for 1 hour. Detected by western blot using chemiluminescence.



EB05062 (3.8 $\mu$ g/ml) staining of paraffin embedded Human Cerebral Cortex. Steamed antigen retrieval with citrate buffer pH 6, AP-staining.