



## UK Office

### Everest Biotech Ltd

Cherwell Innovation Centre  
77 Heyford Park  
Upper Heyford  
Oxfordshire  
OX25 5HD  
UK

Enquiries:

[info@everestbiotech.com](mailto:info@everestbiotech.com)

Sales:

[sales@everestbiotech.com](mailto:sales@everestbiotech.com)

Tech support:

[support@everestbiotech.com](mailto:support@everestbiotech.com)

Tel: +44 (0)1869 238326

[www.everestbiotech.com](http://www.everestbiotech.com)

**Research Use Only. Not for  
diagnostic or therapeutic use.**

## EB09195 - Goat Anti-XCL1 / XCL2 Antibody

Size: 100µg specific antibody in 200µl



### Target Protein

**Principal Names:** XCL1, chemokine (C motif) ligand 1, ATAC, LPTN, LTN, SCM-1, SCM-1a, SCM1, SCYC1, OTTHUMP00000032498, OTTHUMP00000060404, lymphotactin, region containing small inducible cytokine subfamily C, member 1 (lymphotactin), small inducible cytokine subfamily C, member 1 (lymphotactin), small inducible cytokine subfamily C, member 1 (lymphotactin), small inducible cytokine subfamily C, member 2, small inducible cytokine subfamily C, member 1 (lymphotactin)

**Official Symbol:** XCL1 / XCL2

**Accession Number(s):** NP\_002986.1; NP\_003166.1

**Human GeneID(s):** [6375](#)

**Important Comments:** XCL1 and CXL2 are almost identical and therefore this antibody is expected to recognize both proteins.

### Immunogen

Peptide with sequence C-RDVVRSMRKSNTNRN, from the internal region of the protein sequence according to NP\_002986.1; NP\_003166.1.

Please note the [peptide](#) is available for sale.

### Purification and Storage

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

### Applications Tested

**Peptide ELISA:** antibody detection limit dilution 1:16000.

**Western blot:** Preliminary experiments gave an approx 80kDa band in Human Peripheral Blood Mononucleocytes lysates after 0.3µg/ml antibody staining. Please note that currently we cannot find an explanation in the literature for the band we observe given the calculated size of 12.5kDa according to NP\_002986.1 and 12.6kDa according to 003166.1. The 80kDa band was successfully blocked by incubation with the immunizing peptide. We would appreciate any feedback from people in the field - have any results been reported with other antibodies/lysates? Have any further splice variants/modified forms been reported?

### Species Reactivity

**Tested:**

**Expected from sequence similarity:** Human