



## UK Office

### Everest Biotech Ltd

Cherwell Innovation Centre  
77 Heyford Park  
Upper Heyford  
Oxfordshire  
OX25 5HD  
UK

Enquiries:

[info@everestbiotech.com](mailto:info@everestbiotech.com)

Sales:

[sales@everestbiotech.com](mailto:sales@everestbiotech.com)

Tech support:

[support@everestbiotech.com](mailto:support@everestbiotech.com)

Tel: +44 (0)1869 238326

[www.everestbiotech.com](http://www.everestbiotech.com)

**Research Use Only. Not for  
diagnostic or therapeutic use.**

## EB07919 - Goat Anti-YAP1 Antibody

Size: 100µg specific antibody in 200µl



### Target Protein

**Principal Names:** Yes-associated protein 1, 65kDa, YAP, YAP2, YAP65, Yes-associated protein 1, 65 kD, Yes-associated protein 1, 65 kDa, yes-associated protein 2

**Official Symbol:** YAP1

**Accession Number(s):** NP\_006097.1

**Human GeneID(s):** [10413](#)

**Non-Human GeneID(s):** 22601 (mouse), 363014 (rat)

### Immunogen

Peptide with sequence C-TYHSRDESTDS, from the internal region of the protein sequence according to NP\_006097.1.

Please note the [peptide](#) is available for sale.

### Purification and Storage

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

### Applications Tested

**Peptide ELISA:** antibody detection limit dilution 1:8000.

**Western blot:** Approx 70kDa band observed in lysates of cell line MDCK (calculated MW of 50.3kDa according to Dog XP\_536601.2). Data obtained from an anonymous customer. Recommended concentration: 0.5-2µg/ml.

**Additional validation:** This antibody has been successfully used in the following paper: Sikorski et al. (2018) PMID: 30377371.

### Species Reactivity

**Tested:** Dog

**Expected from sequence similarity:** Human, Mouse, Rat, Dog

### Specific Reference


**This antibody has been successfully used in the following paper:**

Krzysztof Sikorski, Adi Mehta, Marit Inngjerdigen, Flourina Thakor, Simon Kling, Tomas Kalina, Tuula A. Nyman, Maria Ekman Stensland, Wei Zhou, Gustavo A. De Souza, Lars Holden, Jan Stuchly, Markus Templin and Fridtjof Lund-Johansen

A high-throughput pipeline for validation of antibodies

Nat Methods. 2018 Nov;15(11):909-912

PMID: 30377371



EB07919 (0.5 $\mu$ g/ml) staining of MDCK lysate (35 $\mu$ g protein in RIPA buffer). Primary incubation was 1 hour.  
Detected by chemiluminescence.