

International Office

Everest Biotech Ltd

Vector Laboratories, Inc. 6737 Mowry Ave Newark, CA 94560 United States

Customer Service:

customerservice@vectorlabs.com

Technical Service:

technical@vectorlabs.com

Tel: +1 (800) 227-6666

www.everestbiotech.com

Research Use Only. Not for diagnostic or therapeutic use.

EB11188 - Goat Anti-ZNF652 (aa594-606) Antibody

Size: 100µg specific antibody in 200µl



Target Protein

Principal Names: DKFZp781E2122, KIAA0924, OTTHUMP00000220047, zinc finger

protein 652, ZNF652 **Official Symbol:** ZNF652

Accession Number(s): NP_055712.1

Human GenelD(s): 22834

Important Comments: Reported variants represent identical protein: NP_055712.1,

NP_001138837.1

Immunogen

Peptide with sequence C-RHLAEKNSSAQHH, from the C Terminus of the protein sequence according to NP_055712.1.

Please note the peptide is available for sale.

Purification and Storage

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

Applications Tested

Peptide ELISA: antibody detection limit dilution 1:64000.

Western blot: Approx 90kDa band observed in Human Heart, Kidney and Liver lysates (calculated MW of 69.7kDa according to NP_055712.1). The observed molecular weight corresponds to earlier findings in literature with different antibodies (Kumar et al, Mol Cancer Res. 2006 Sep;4(9):655-65.PMID: 1696643). Recommended concentration: 0.01-0.03µg/ml. Primary incubation was 1 hour.

IHC: Paraffin embedded Human Small Intestine. Recommended concentration: 5µg/ml.

Species Reactivity


Tested: Human

Expected from sequence similarity: Human, Cow, Pig



20kDa

EB11188 (0.01µg/ml) staining of Human Heart lysate (35µg protein in RIPA buffer). Detected by chemiluminescence.



 $EB11188 \ (5\mu g/ml) \ staining \ of \ paraffin \ embedded \ Human \ Small \ Intestine. \ Steamed \ antigen \ retrieval \ with \ citrate$ buffer pH 6, AP-staining.