

#### **UK Office**

**Everest Biotech Ltd** 

Cherwell Innovation Centre

77 Heyford Park Upper Heyford Oxfordshire

OX25 5HD UK

Enquiries:

info@everestbiotech.com

Sales:

sales@everestbiotech.com

Tech support:

support@everestbiotech.com

Tel: +44 (0)1869 238326

www.everestbiotech.com

Research Use Only. Not for diagnostic or therapeutic use.

# EB09206 - Goat Anti-ZO-1 Antibody

Size: 100µg specific antibody in 200µl



# **Target Protein**

**Principal Names:** TJP1, tight junction protein 1 (zona occludens 1), DKFZp686M05161, MGC133289, ZO-1, OTTHUMP00000174520, tight junction protein 1, tight junction

protein ZO-1, zona occludens 1, zonula occludens 1 protein

Official Symbol: TJP1

Accession Number(s): NP\_003248.3; NP\_783297.2

Human GeneID(s): 7082

Important Comments: This antibody is expected to recognize both reported isoforms

(NP\_003248.3; NP\_783297.2)

## Immunogen

Peptide with sequence C-PKPTSQNQFSEHDKT, from the internal region of the protein sequence according to NP\_003248.3; NP\_783297.2.

Please note the peptide is available for sale.

### **Purification and Storage**

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin

Aliquot and store at -20°C. Minimize freezing and thawing.

## **Applications Tested**


Peptide ELISA: antibody detection limit dilution 1:64000.

**Immunofluorescence:** Strong expression of the protein seen in the membrane, cytoplasm and nuclei of U251 cells. Recommended concentration: 10µg/ml.

#### **Species Reactivity**

Tested: Human

Expected from sequence similarity: Human



EB09206 Immunofluorescence analysis of paraformaldehyde fixed U251 cells, permeabilized with 0.15% Triton. Primary incubation 1hr (10ug/ml) followed by Alexa Fluor 488 secondary antibody (2ug/ml), showing membrane, cytoplasmic and some nuclear staining. The nuclear stain is DAPI (blue). Negative control: Unimmunized goat IgG (10ug/ml) followed by Alexa Fluor 488 secondary antibody (2ug/ml).